Catherine J Frompovich 23 Cavendish Drive, Ambler, PA 19002 215-653-7575

June 7, 2017

## Certification of Frompovich Exceptions Protest June 7, 2017 Posted U.S. Postal Service June 7, 2017

Catherine J Frompovich v. PECO Energy Company, C-2015-2474602

<u>Certified Mail \* Return Receipt Requested</u> Rosemarie Chiavetta, Secretary Pennsylvania Public Utility Commission Commonwealth of Pennsylvania P.O. Box 3265 Harrisburg, PA 17105-3265

Courtesy copy to OSA emailed to ra-OSA@pa.gov

First Class USPS Mail with Tracking copies to: Administrative Law Judge Darlene D Heep Pennsylvania Public Utility Commission 801 Market Street, Ste. 4063 Philadelphia, PA 19107

Ward Smith, Esq. Shawane Lee, Esq. Thomas Carl Watson, Esq. Exelon Business Services Company LLC Legal Department 2301 Market Street, S23-1 Philadelphia, PA 19103

Respectfully submitted by,

atturing Tromporick

Catherine J'Frompovich, Pro Se

# RECEIVED

JUN 7 2017

PA PUBLIC UTILITY COMMISSION SECRETARY'S BUREAU

## Case No. C-2015-2474602 PENNSYLVANIA PUBLIC UTILILTY COMMISSION

CATHERINE J. FROMPOVICH

vs.

PECO ENERGY COMPANY

## CATHERINE J. FROMPOVICH EXCEPTIONS PROTEST OF INITIAL DECISION

# RECEIVED

JUN -7 2017

PA PUBLIC UTILITY COMMISSION SECRETARY'S BUREAU

Catherine J. Frompovich, Pro Se 23 Cavendish Drive Ambler, PA 19002 215-653-7575

# BEFORE THE FENNSYLVANIA PUBLIC UTILITY COMMISSION

;

RECEIVED

JUN -7 2017

Catherine J. Frompovich v. PA PUBLIC UTILITY COMMISSION SECRETARY'S BUREAU

C-2015-2474602

PECO Energy Company

## EXCEPTIONS PROTEST OF INITIAL DECISION

1. Catherine J. Frompovich, an almost 79-year-old breast cancer survivor, respectfully files a protest rather than an appeal, which would be pointless especially since the PA PUC's Administrative Law Court is nothing short of an apparent legal appendage of the PA Public Utility Commission. Said PA PUC apparently is not interested in protecting the public from unsafe utility innovations and/or consensus microwave industry associations pseudoscience spouted as 'gospel truth' by vested microwave industry trade associations and further parroted by utility company expert witnesses, who are regarded as the only source of scientific information.

2. One environmentally risky example of PECO's unsafe utility innovations is recommending and encouraging the use of CFL (compact fluorescent lamp) light bulbs which, according to ¶1 on page 22 of the Initial Decision "...florescent lights have particularly strong harmonics..." which emit "dirty electricity" into homes and wherever CFLs are used. Plus, CFLs contain mercury, which creates a hazmat situation when CFLs are broken, plus they normally are problematic to dispose of.

3. Frompovich presented hundreds, if not thousands, of document citations and/or actual documents at various times during the course of these protracted proceedings regarding adverse health effects from non-thermal radiation waves emitted by microwave technologies, which

operate utility company AMI Smart Meters, only to be totally disregarded and probably procedurally discarded, despite 32 percent of microwave-industry-funded research studies indicating adverse effects occur. That fact alone should negate the PA PUC's, PECO's and its expert witnesses', and this Court's close-minded opinions and acceptance of non-existence of non-thermal radiation wave adverse effects upon humans, animals and vegetation. Non-thermal waves "non-existence" is promoted by one of the microwave industry's key professional associations, ICNIRP, which has been cited for conflicts of interest and other questionable citations as reported in Frompovich's January 20, 2017 Brief *cf. Brief 40-45 (102-115)*.

4. Frompovich's attempted introductions during the November 2-3, 2016 hearing of published peer review studies regarding cancer and EMF/RF/ELF, *specifically breast cancers*, were rejected and not permitted to be entered into the hearing record; something this Court was remiss in by sustaining PECO's objections time after time.

5. Administrative Law Courts often are referred to as "captive" and "kangaroo courts" with their status being questioned.

"In her ruling, Judge [Leigh Martin] May attacked an important element of the system of independent agencies first set up during the administration of President Franklin Delano Roosevelt, under which Congress delegates its authority to administer complex regulations to federal agencies overseen by supposedly independent boards. Those agencies also can use administrative law judges, operating outside the rules requiring life tenure and other protections under Article III of the Constitution, to decide civil complaints."<sup>1</sup>-<sup>2</sup>

"This new decision calling the Administrative Law Courts what they really are is reminiscent of the notorious extrajudicial proceedings of the Star Chamber operated by King James I. The court of Chancery set up outside of the King's Bench, so there were no trials by jury. It had the same purpose, to circumvent the law. This is where our Fifth

<sup>&</sup>lt;sup>1</sup> <u>https://www.forbes.com/sites/danielfisher/2015/06/09/did-a-judge-just-kick-off-the-great-unraveling-of-the-administrative-state-with-sec-ruling/#4572610e50ac</u> accessed 6-5-17

<sup>&</sup>lt;sup>2</sup> <u>https://www.scribd.com/document/268154852/Hill-v-SEC-AU-Order\_accessed 6-5-17</u>

Amendment privilege came into being. That came about following the trial of John Lilburne (1615-1657) for handing out a pamphlet the government did not like."<sup>3</sup>

"Not discussed in the coverage of this story is that the Administrative Law Courts are a fiefdom, to put it mildly. They have long been corrupt and traditionally rule in favor of their agencies, making it very costly for anyone to even try to defend themselves. If someone were to attempt this feat, first they have to wear the costs of an Administration proceeding and appeal to an Article III court judge, then they must appeal to the Court of Appeals, and finally plea to the Supreme Court. The cost of such adventures is well into the millions, and good luck on actually getting justice."<sup>4</sup>

### Furthermore,

"One major difference between ALJs and traditional judges is that ALJs serve as both the judge and trier of fact. This is known as a bench trial. In civil court, the parties sometimes have the option of choosing to forgo a jury and have the judge weigh the factual evidence that the parties provide. During an administrative hearing, however, the ALJ will always weigh the evidence and make factual determinations."<sup>5</sup>

The book *Is Administrative Law Unlawful?* "reveals administrative law to be not a benign, natural outgrowth of contemporary government but a pernicious—and profoundly unlawful—return to dangerous pre-constitutional absolutism."<sup>6</sup> The author is Philip Hamburger, the Maurice and Hilda Friedman Professor of Law at Columbia Law School.

Some critics of administrative law contend it, undoubtedly, is stacked in favor of government.

6. Frompovich alleges ALJ Darlene D. Heep not only did not weigh the evidence in Frompovich's case, Judge Heep actively prevented evidence, which was germane to Frompovich's case, e.g., cancer studies and EMFs/RFs/ELFs, from being introduced during the hearing, at the behest and objections from PECO's attorneys; said objections were sustained for most of the exhibits Frompovich tried to introduce into the record. Frompovich, therefore, feels and also alleges she did not receive a fair and judicious hearing. Frompovich further contends

<sup>&</sup>lt;sup>3</sup> <u>https://www.armstrongeconomics.com/history/americas-economic-history/judge-rules-administrative-court-</u> <u>system-illegal-after-81-years/</u> accessed 6-5-17

<sup>&</sup>lt;sup>4</sup> Ibid.

<sup>&</sup>lt;sup>5</sup> <u>https://www.justia.com/administrative-law/administrative-law-judges/</u> accessed 6-6-17

<sup>&</sup>lt;sup>6</sup> <u>http://press.uchicago.edu/ucp/books/book/chicago/I/bo17436684.html</u> accessed 6-6-17

she was discriminated against insofar as Judge Heep's Initial Decision *Discussion* totally ignores the Americans with Disabilities *Amendments Act*, which grants Frompovich special health considerations, which this Court apparently deliberately ignores, but Judge Heep should know better, since she's a known expert in ADA law.

7. Frompovich alleges Judge Heep probably did not write much of the Initial Decision, as

it seemingly has PECO attorney Ward Smith's 'fingerprint' writing style and mistakes contained

within.

2

One specific example of that is found on page 18:

"Ms. Frompovich contends that installation of a Smart Meter at her home would counter, and negatively affect, her healing and health. She also asserts that she is protected by, and the PECO is subject to, the Americans with Disabilities Act (ADA), 42 U.S.C.S. § 12132, et seq. because she had cancer."

Throughout all of Frompovich's filings and testimony she specifically cited the

Americans with Disabilities Act Amendments Act<sup>7</sup> §1630.2 (G) Disability; §1630.2 (J) Substantially Limits; §1630.2 (I) Major Life Activities; §1630.2(j)(1)(ii) Significant or Severe Restriction Not Required Nonetheless, Not Every Impairment Is Substantially Limiting; §1630.2(j)(1)(iii) Substantial Limitation Should Not Be Primary Object of Attention; Extensive Analysis Not Needed. (cf. Brief pp. 14-16)

PECO Attorney Ward in both his Brief and Respondent Brief totally neglects addressing the

ADA Amendments Act, which also would impact PECO, but prefers only to cite the original

ADA rather than the updated health considerations and benefits Congress passed as law in the

Amendments Act<sup>8</sup>.

8. It is Frompovich's contention AL Judge Heep could not have omitted the importance

and very essence of the ADA Amendments Act affecting cancer patients because AL Judge Heep

is considered an expert in the Americans with Disabilities Act, which also includes the later add-

<sup>&</sup>lt;sup>7</sup> Pub.L. 110-325 ADAAA

<sup>&</sup>lt;sup>8</sup> Ibid

on Amendments Act. AL Judge Heep apparently presided over a 2015 hearing regarding the Rhode Island Disability Law Center in Providence, Rhode Island.

9. Another example, which could indicate Attorney Ward's sloppy legal writing or his contribution thereto, is Ward's apparent proclivity for misstatement of facts, as Frompovich pointed out numerous times in her Response Brief dated February 10, 2017. In the current Initial Decision, Frompovich's book is cited on page 6 as "The Cancer Answer," which she did not write; several other authors wrote books with such titles. For the record, Frompovich wrote "A

## Cancer Answer, Holistic BREAST Cancer Management."

10. Frompovich wishes to state categorically the Initial Decision is replete with misstatements and downright falsehoods! Some examples are Clause 53, page 10; Clause 60, page 11, which can be clarified by PECO's expert witness Glenn Pritchard as to **the facts** when Mr. Pritchard answered AL Judge Christopher Pell's questioning about smart meters *cf. Transcript Pg. 169 (1-5)* when Mr. Pritchard said, "It could be once every five minutes to once every hour or maybe once a day depending on what the device – whether it would be a smart thermostat, a dishwasher as you mentioned or maybe an in-home display devise."

Quoting from Frompovich's Respondent Brief Clause 24, Page 17, here are the facts:

"Judge Pell then remarked, 'I understand you to say that, if it doesn't connect with anything, it pulses every 30 seconds?" PECO employee and expert Pritchard states, 'It continues to seek that, yes.' Whereas, His Honor Judge Pell then asks, 'Indefinitely or will it decide, okay, I'm not finding anything, stop? Can that be adjusted?' Mr. Pritchard replied, 'No.' Judge Pell queries further, 'No. Does it have to be that way?' Mr. Pritchard replied, 'We have no options with that.' cf. Transcript Pg. 169 (1-17)"

11. Frompovich alleges AL Judge Heep could not have made such an obvious mistake in writing Clause 60 unless she is in PECO's back pocket, which Frompovich seriously doubts AL Judge Heep is. Clause 60 is wordsmithed to protect PECO's *Focus* AMI Smart Meter's almost

5

# PECO's Multi-Tiered Smart Grid Network



## Communication Tiers:

- Tier-1: ~375 miles of high speed/high bandwidth fiber optic communications
- ✓ Tier-2: WiMax wireless communication network
- Tier-3: AMI Network, low-speed, low bandwidth network
- ✓ Tier-4: HAN In-Home Communications



constant search-for-appliance-or-device microwave GHz transmissions as **Mr. Pritchard's remarks quoted above prove,** and is a blatant misrepresentation in the Initial Decision. How **can the PA PUC and this Court accept apparent LIES as facts, truths and science in the Initial Decision?** Can you please answer that, Judge Heep, as you signed the document? During the November hearing, PECO constantly referred to their smart meter as a *FlexNet* AMI Smart Meter. It's now a *Focus* meter?

12. Clauses 61 and 62 are at variance with what utilities are doing regarding collection and selling of data as described by the 2 minute video explaining the algorithm "Onzo"<sup>9</sup> used in Smart Meter data mining.

13. On page 21 of the Initial Decision, ¶2 discusses PECO's AMI meters wherein "He [Mr. Pritchard] stated that while some other utilities employ a mesh system, which transfers data from meter to meter until it reaches a data collector, resulting in many transmissions, in the PECO system, messages are transmitted directly to the collector and do not have to transmit from one meter to the next in a relay, resulting in fewer transmissions."

14. Frompovich points out again, as she did in her Respondent's Brief "Exhibit No. 4— Glenn Pritchard, PE, IEEE Presentation, PECO delivers a Reliable and Resilient Smart Grid PowerPoint Frame 8: PECO's Multi-Tiered Smart Grid Network," and now includes a photocopy of that Multi-Tiered Smart Grid NETWORK. What is that four-tier network all about?

15. Another factual anomaly appears on page 23 where "Dr. Christopher Davis, PhD, testified that, to a reasonable degree of scientific certainty, 'AMI meters are incapable of causing any biological effects, certainly no adverse biological effects in anybody.' (Tr. 216)"

:

<sup>&</sup>lt;sup>9</sup> <u>https://youtu.be/uluKjzqHDz0?t=8</u> accessed 6-5-17

Dr. Davis' "wiggle-room" statement is patently incorrect and factually challenged by two videos showing the effects of individuals hooked up to EKGs that register the PQRST waves of the human heart when an AMI Smart Meter is activated during a test monitored by a medical doctor. Both tests and their results can be read and seen in two of my recent *Activist Post* articles:

Can—And Do—Utility Smart Meters Affect The Human Heart http://www.activistpost.com/2017/06/can-utility-smart-meters-affect-human-heart.html



AMI Smart Meters Interfere With Human Heart Rhythms, EGK Proves http://www.activistpost.com/2017/05/ami-smart-meters-interfere-human-heart-rhythms-ekgproves.html

16. Illustrating, plus further compounding Dr. Christopher Davis, PhD, and Dr. Mark A. Israel, MD, both PECO EMF experts of sorts testifying in the Frompovich matter, unquestionable apparent lack of current scientific information regarding microwave EMFs, Frompovich introduces **Exhibit 1**, the August 2015 *Journal of Chemical Neuroanatomy* article "Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric

effects including depression," Professor Emeritus of Biochemistry and Basic Medical Sciences,

Washington State University, Martin L. Pall published indicating:

2

"1. Such EMFs act via activation of VGCCs [voltage-gated calcium channels], acting through the VGCC voltage sensor which is predicted to be exquisitely sensitive to these EMFs (Pall, 2015). VGCCs occur in high densities throughout the nervous system and have essential roles throughout the nervous system in releasing neurotransmitters and neuroendocrine hormones.

"2. Elevated EGCC activity, produced by an allete of the CACNAIC gene which encodes the channel of the main L-type VGCC in the brain, produces various neuropsychiatric effects (Table 1). This predicts, that low intensity non-thermal microwave frequency EMFs which also produce elevated L-type and other VGCC activity, therefore produce widespread neuropsychiatric effects."

"3. Studies reviewed in the Tolgskaya and Gordon, 1973 publication (Table 2) have shown that the cells of the mammalian nervous system show high sensitivity to various non-thermal microwave and lower frequency EMFs, being, apparently more sensitive than any other organ in the body of the rodents. These studies predict that the human nervous system is likely to be similarly sensitive to these EMFs, predicting, therefore, widespread neuropsychiatric effects in humans."

Based upon the above science, it is conceivable the PA PUC and all utility companies mandating

retrofitting AMI Smart Meters emitting non-thermal radiation waves are opening themselves to

class action lawsuits, which should be instituted as soon as possible, especially since there are

increasingly escalating erratic behavioral and mental health issues within the USA.

According to the U.S. National Institute of Mental Health,

- Mental and behavioral disorders, which account for 13.6 percent of total U.S. DALYs [disability-adjusted life years]; and,
- Neurological disorders, which account for 5.1 percent of total U.S. DALYs<sup>10</sup>

What EHS-, mental-, and behavioral-related issues are AMI Smart Meters contributing to the above statistics since people in Pennsylvania are mandated to live with non-thermal radiation

<sup>&</sup>lt;sup>10</sup> <u>https://www.nimh.nih.gov/health/statistics/disability/us-leading-categories-of-diseases-disorders.shtml</u> accessed 6-6-17

waves and adverse effects 24/7/365? The PA PUC certainly should be ashamed of its cavalier attitudes and administrative regulations in not protecting the health and wellbeing of Pennsylvania citizens, especially growing children who are impacted more seriously by EMFs/RFs/ELFs.



Slides from Dr. Om Ghandi, As cited by the World Health Organization's International Agency for the Research on Cancer

Source: Google Charts

Ignorance, therefore, is not a PA PUC defense, since numerous Pennsylvanians have protested and brought attention to their health problems known as electromagnetic hypersensitivity (EHS) from EMFs/RFs/ELFs, which AMI Smart Meters emit, and the PA PUC deliberately ignores.

17. Additionally, the *Bioinitiative Report* (which Frompovich introduced as Exhibit 2 on PECO Cross-examination) has documented thousands of adverse health studies from non-thermal radiation waves from EMFs/RFs/ELFs, which ICNIRP and industry-consensus-science

deny as possible or even "exist." Certainly, if industry-funded research found 32 percent of studies confirm such effects, how in the name of sanity and science-based medicine can they be disregarded as non-existent? Is there some sort of collusion to deceive or achieve a special agenda? The very fact almost one-third (32%) found non-thermal effects proves they do exist! PECO, the PA PUC and even this Court seem to be acting in some sort of complicity for, or to effectuate, a specific purpose which, apparently, is not in the best health interests, safety and wellbeing of Pennsylvania utility customers, Claimant Frompovich included.

18. There are many more issues in the Initial Decision Frompovich could discuss and provide scientific evidence pro and con, but what's the use, since there apparently is a set agenda in Pennsylvania regarding AMI Smart Meters and the illegal problems they apparently are creating for consumers, their health and otherwise, e.g., meter fires, exorbitant *Time of Use* rates, surveillance, and customer-data-mining by utilities to sell for marketing purposes<sup>11</sup>.

19. Frompovich respectfully leaves all responsible parties involved in the AMI Smart Meter legal, health and political debacle in Pennsylvania with these words of wisdom:

Isaiah 10

Woe to those who enact unjust statutes and who write oppressive decrees Depriving the needy of judgment and robbing my people's poor of their rights... Source: The International Student Bible for Catholics, Thomas Nelson Publishers

Proverbs 1:7 To have knowledge, you must first have reverence for the Lord. Stupid people have no respect for wisdom and refuse to learn. Source: Good News Bible, American Bible Society

Respectfully submitted by

Jumporite

Catherine J. Frompovich, Pro Se

June 7, 2017

<sup>&</sup>lt;sup>11</sup> https://youtu.be/uluKjzgHDz0?t=8 accessed 6-5-17



## JUN -7 2017

PA PUBLIC UTILITY COMMISSION SECRETARY'S BUREAU

.

## **EXHIBIT 1**

.

## JOURNAL OF CHEMICAL NEUROANATOMY

Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression

August 2015

Contents lists available at ScienceDirect



Review

Journal of Chemical Neuroanatomy



journal homepage: www.elsevier.com/locate/jchemneu

## Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression



## Martin L. Pall

Professor Emeritus of Biochemistry and Basic Medical Sciences, Washington State University, 638 NE 41st Avenue, Portland, OR 97232-3312, USA

#### ARTICLE INFO

Article history: Received 13 April 2015 Received in revised form 1 August 2015 Accepted 9 August 2015 Available online 21 August 2015

Keywords: Excessive calcium effects Oxidative/nitrosative stress Low-intensity microwave electromagnetic fields

### ABSTRACT

Non-thermal microwave/lower frequency electromagnetic fields (EMFs) act via voltage-gated calcium channel (VGCC) activation. Calcium channel blockers block EMF effects and several types of additional evidence confirm this mechanism. Low intensity microwave EMFs have been proposed to produce neuropsychiatric effects, sometimes called microwave syndrome, and the focus of this review is whether these are indeed well documented and consistent with the known mechanism(s) of action of such EMFs. VGCCs occur in very high densities throughout the nervous system and have near universal roles in release of neurotransmitters and neuroendocrine hormones. Soviet and Western literature shows that much of the impact of non-thermal microwave exposures in experimental animals occurs in the brain and peripheral nervous system, such that nervous system histology and function show diverse and substantial changes. These may be generated through roles of VGCC activation, producing excessive neurotransmitter/neuroendocrine release as well as oxidative/nitrosative stress and other responses. Excessive VGCC activity has been shown from genetic polymorphism studies to have roles in producing neuropsychiatric changes in humans. Two U.S. government reports from the 1970s to 1980s provide evidence for many neuropsychiatric effects of non-thermal microwave EMFs, based on occupational exposure studies. 18 more recent epidemiological studies, provide substantial evidence that microwave EMFs from cell/mobile phone base stations, excessive cell/mobile phone usage and from wireless smart meters can each produce similar patterns of neuropsychiatric effects, with several of these studies showing clear dose-response relationships. Lesser evidence from 6 additional studies suggests that short wave, radio station, occupational and digital TV antenna exposures may produce similar neuropsychiatric effects. Among the more commonly reported changes are sleep disturbance/insomnia, headache, depression/depressive symptoms, fatigue/tiredness, dysesthesia, concentration/attention dysfunction, memory changes, dizziness, irritability, loss of appetite/body weight, restlessness/anxiety, nausea, skin burning/tingling/dermographism and EEG changes. In summary, then, the mechanism of action of microwave EMFs, the role of the VGCCs in the brain, the impact of non-thermal EMFs on the brain, extensive epidemiological studies performed over the past 50 years, and five criteria testing for causality. all collectively show that various non-thermal microwave EMF exposures produce diverse neuropsychiatric effects.

© 2015 The Author. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

#### Contents

1.	Introduction	44
2.	Microwave/lower frequency EMFs act to activate voltage-gated calcium channels	44
3.	Genetic polymorphism studies	44
4.	Histological and functional changes in central nervous system (CNS) and peripheral nervous system (PNS) in animals exposed	
	to microwave EMFs	44
5.	Older epidemiological reviews and other related studies	45

http://dx.doi.org/10.1016/j.jchemneu.2015.08.001

0891-0618/@ 2015 The Author. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

E-mail address: martin\_pall@wsu.edu.

6.	Specific epidemiological studies on neuropsychiatric effects of microwave EMFs	45
7.	Criteria for assessing causality in epidemiological studies	46
8.	Discussion and conclusions	48
	References	50

Chemicals having roles:

Calcium(2+) Nitric oxide (NO) Oxido nitrite (peroxynitrite)

#### 1. Introduction

Microwave syndrome (Hocking, 2001; Johnson Liakouris, 1998), a combination of various neuropsychiatric symptoms originally described in persons with occupational exposures to microwave frequency EMFs, has been disputed largely because of the lack of an apparent mechanism for generating these symptoms. It is reported to often include such symptoms as fatigue, headache, insomnia, dysesthesia (impaired sensation), irritability, lack of concentration and other symptoms (Hocking, 2001; Johnson Liakouris, 1998). Similar but more extensive combinations of symptoms have been reported following occupational exposures in two U.S. government reports from the 1970s/1980s (Naval Medical Research Institute Research Report, 1971; Raines, 1981) and following environmental exposures as described in two more recent reviews (Khurana et al., 2010; Levitt and Lai, 2010).

The goal here is not just to review the epidemiology, however, but more importantly to consider the issue of possible physiological mechanism(s). Hennekens and Buring (1989), on p. 40 in their textbook Epidemiology in Medicine state "The belief in the existence of a cause and effect relationship is enhanced if there is a known or postulated biologic mechanism by which the exposure might reasonably alter risk of developing disease." It is of critical importance therefore to assess possible biological mechanism before considering the epidemiological evidence.

Accordingly, this paper considers the mechanism by which low intensity microwave EMFs impact the cells of our bodies, how that mechanism may be predicted to impact the nervous system, evidence for such impact from experimental animal studies, genetic polymorphism evidence for that mechanism acting in humans to produce neuropsychiatric effects and finally, the epidemiological evidence for such effects in human populations with repeated low level microwave EMF exposure. Consideration of each of these types of evidence influences the overall interpretation presented in this paper.

#### 2. Microwave/lower frequency EMFs act to activate voltagegated calcium channels

In 24 different studies reviewed earlier (Pall, 2013) and two additional studies (Li et al., 2014; Lisi et al., 2006), microwave and lower frequency low intensity EMF effects were blocked or greatly lowered by calcium channel blockers, agents thought to be specific for blocking voltage-gated calcium channels (VGCCs). In these 26 studies, a total of 5 distinct types of channel blockers were used, with each type having a distinct structure and binding to a distinct site, such that it is essentially certain that these must be acting by blocking VGCCs, which is their only known common property. In each of these 26 studies, each of the responses studied, were blocked or greatly lowered by calcium channel blockers, showing that VGCC activation has roles in producing a wide variety of EMF effects. There is a large literature on changes in calcium fluxes and in calcium signaling following microwave EMF exposure (partially reviewed in Walleczek, 1992; Adey, 1993); each of these, including calcium efflux changes, can be explained as being due to VGCC activation, again suggesting a widespread role of VGCC activation in producing biological responses to EMFs. Pilla (2012) showed that pulsed microwave field exposure, produced an almost instantaneous increase in calcium/calmodulin-dependent nitric oxide (NO) signaling, providing strong evidence that these fields can produce an almost instantaneous VGCC activation. It is likely, that these EMFs act directly on the voltage sensor of the VGCCs to produce VGCC activation (Pall, 2015) with the voltage sensor being exquisitely sensitive to these EMFs because of its physical properties and location in the plasma membrane.

EMFs have been proposed to act to produce a wide variety of responses in the cell, via downstream effects of VGCC activation (Pall, 2013, 2014, 2015), including elevated intracellular calcium [Ca2+]i, excessive calcium and nitric oxide signaling and also excessive peroxynitrite, free radicals and oxidative stress.

VGCC activation has been shown to have a universal or nearuniversal role in the release of neurotransmitters in the brain and also in the release of hormones by neuroendocrine cells (Berridge, 1998; Dunlap et al., 1995; Wheeler et al., 1994), with such release being produced by calcium signaling. There are high densities of diverse VGCCs occurring in neurons throughout the nervous system. Both the high VGCC density and their function in neurotransmitter and neuroendocrine release throughout the nervous system suggests that the nervous system is likely to be highly sensitive to low intensity EMFs.

#### 3. Genetic polymorphism studies

Genetic polymorphism studies are powerful tools for looking at the roles of specific proteins in human populations. In Table 1, a series of genetic polymorphism studies have been performed that show that an allele producing increased expression of the gene encoding the channel of the main L-type VGCC in the brain, produces diverse neuropsychiatric effects. These studies clearly show that excess L-type VGCC activity can cause neuropsychiatric effects. They also predict, therefore, that increased VGCC activity produced by microwave EMFs may be able to also produce widespread neuropsychiatric effects.

# 4. Histological and functional changes in central nervous system (CNS) and peripheral nervous system (PNS) in animals exposed to microwave EMFs

The most extensive literature on histological and functional changes in animals is from the Soviet literature from the 1950s/ 1960s with additional Western literature from the same time period. Both Soviet and non-Soviet literature were reviewed in an English language Publication by Tolgskaya and Gordon (1973). This publication is, therefore, the main focus of this section. That publication was divided into thermal and non-thermal exposure studies, with the non-thermal studies which occupy the majority of the text (pp. 53–137) being of sole interest here.

#### Table 1

Influence of genetic polymorphism of the CACNA1C in producing diverse neuropsychiatric effects.

Citation	Genetic polymorphism	Changes produced by allele of gene
Bhat et al. (2012)	Polymorphism producing Increased expression of CACNATC L-type VGCC subunit	Review: The polymorphism Is associated with increased susceptibility to bipolar disorder, "depression, schizophrenia, autism spectrum disorders, as well as changes in brain function and structure in control subjects who have no diagnosable psychiatric illness."
Bigos et al. (2010)	Polymorphism producing Increased expression of CACNA1C L-type VGCC subunit	Associated with increases in both bipolar disorder and schizophrenia
Krug et al. (2010)	Polymorphism producing increased expression of CACNA1C L-type VGCC subunit	Negatively influences language production on a semantic level
Krug et al. (2014)	Polymorphism producing increased expression of CACNA1C L-type VGCC subunit	Influences episodic memory and retrieval
Soeiro-de-Souza et al. (2012)	Polymorphism producing increased expression of CACNA1C L-type VGCC subunit	Produces impaired facial emotion recognition
Tesli et al. (2013)	Polymorphism producing increased expression of CACNA1C L-type VGCC subunit	Produces increased activation of the amygdala during emotional processing
Thimm et al. (2011)	Polymorphism producing increased expression of CACNA1C L-type VGCC subunit	Associated with attention deficits including alerting, orienting and executive control of attention

These were all derived from the Tolgskaya and Gordon (1973) review and page numbers listed are page numbers from that document. All refer to changes produced by non-thermal exposures in the nervous system of experimental animals, with most being in rats.

This discussion scrolls down through Table 2.

The majority of the histological changes seen in these mostly rodent studies, are seen in the nervous system, despite its being less than 2% of the rodent cell mass. There are statements made that the nervous system, both central and peripheral, is the most highly sensitive tissue to these non-thermal microwave and lower frequency EMFs. Following the nervous system in sensitivity are the myocardium and the testis; myocardial cells are known to have very high densities of VGCCs with especially high densities in the pacemaker cells and the testis is known to have high densities specifically of the T-type VGCCs. Pulsed EMFs are more active in producing histological changes in the brain than are non-pulsed fields, in two studies reviewed; there is a much larger literature showing that in most cases pulsed fields are more biologically active (Pall, 2015; Pangopoulos et al., 2013; Belyaev, 2015).

A wide variety of brain and peripheral nervous system tissues show histological changes following non-thermal exposures. Among the important tissues impacted are the hypothalamus and pituitary gland, where both show similar patterns of changes in neuroendocrine activities. There Is an initial increase in neuroendocrine activity (this may be produced directly by VGCC stimulation of secretion), followed over time by "exhaustion" of neuroendocrine activity (this may be produced by tissue damage produced from long term intracellular calcium [Ca2+]i elevation).

There are widespread histological changes produced in neuronal and neuroendocrine tissues. These were repeatedly reported to be largely reversible on cessation of EMF exposure. They become, however, irreversible when exposure is extended in time. There are changes in EEG activity, which may be an easily measurable monitor of neurological damage.

In a summary statement, Tolgskaya and Gordon (1973) state, "This does not confirm the view, so widely held in the past among Soviet investigators and still maintained to a large extent even at the present time in the West, that the action of microwaves is entirely thermal."

While there were many studies of brain impact of non-thermal EMFs performed in the 1950s/60s that make the information content of Tolgskaya and Gordon (1973) quite high, there is also a substantial recent literature on brain effects of non-thermal microwave EMF exposures (see, for example: Ammari et al., 2008a,b; Bas et al., 2009; Brillaud et al., 2007; Carballo-Quintás et al., 2011; Eberhardt et al., 2008; Dasdag et al., 2009, 2012;

Grafström et al., 2008; Kumlin et al., 2007; López-Martín et al., 2006; Mausset-Bonnefont et al., 2004; Odaci et al., 2008; Rağbetli et al., 2010; Salford et al., 2003; Sonmez et al., 2010).

#### 5. Older epidemiological reviews and other related studies

Two U.S. Government reports each listed many apparent neuropsychiatric effects of microwave/radiofrequency EMFs and a third recognized the role of non-thermal effects on our bodies, but had only a little consideration of neuropsychiatric effects.

The earliest to these was a Naval Medical Research Institute (NMRI) Research Report (1971) which listed 40 apparent neuropsychiatric changes produced by non-thermal exposures including: 5 central/peripheral nervous system (NS) changes, 9 CNS effects, 4 autonomic system effects, 17 psychological disorders, 4 behavioral changes and 2 misc. effects. This NMRI report also provided a supplementary document listing over 2300 citations documenting these and other effects of microwave exposures in humans and in animals.

The Raines (1981) NASA report reviewed extensive literature based on occupational exposures to non-thermal microwave EMFs, with that literature coming from U.S., Western European and Eastern European studies. There are no obvious differences in the literature coming from these different regions. Based on multiple studies, Raines (1981) reports 19 neuropsychiatric effects to be associated with occupational microwave/radiofrequency EMFs.

The Bolen (1994) report put out by the Rome Laboratory of the U.S. Air Force, acknowledged the role of non-thermal effects of microwave EMFs on humans. This report states in the Conclusion section that "Experimental evidence has shown that exposure to low intensity radiation can have a profound effect on biological processes. The nonthermal effects of RF/MW radiation exposure are becoming important measures of biological interaction of EM fields." Clearly Bolen (1994) rejects the claim that only thermal effects occur. Bolen (1994) discusses a specific non-thermal neuropsychiatric effect, where anesthetized animals are awakened when the head is irradiated with microwave EMFs. This suggests a similar mechanism to that acting in humans where such EMFs produce insomnia (see below).

## 6. Specific epidemiological studies on neuropsychiatric effects of microwave EMFs

There are 26 different epidemiological studies described in Table 3. Although 4 of these only studied a single neuropsychiatric effect, 22 of these each provide substantial evidence for the pattern described in the earlier U.S. reports, that a wide range of

#### Table 2

Histological and functional changes in brain function in animals following exposure to non-thermal microwave EMFs.

Observations including page numbers	Comment from Author
The majority of the histological changes seen following non-thermal exposures, occurred in the nervous system, despite its being only about 2% of the tissue mass in rodents; this suggests that the nervous system is highly sensitive to such exposures. Elsewhere (pp. 129, 136), it is suggested that the nervous system is the most sensitive tissue, followed by the heart and the testis, among all of the tissues of the body. The most severe histological changes produced by these non-thermal EMF exposures occur in the nervous system (pp. 136).	High CNS sensitivity to EMFs is predicted by the high density of VGCCs that occur in neurons throughout the nervous system, plus the VGCC role in neurotransmitter and neuroendocrine release.
Pulsed fields were more active than non-pulsed fields in producing histological changes (pp. 71, 97).	Pulsed fields have often been found to be more biologically active than are non-pulsed fields in many different studies from many countries (Pall, 2015; Pangopoulos et al., 2013; Belyaev, 2015).
Nervous system regions impacted by non-thermal microwave and lower frequency fields include: cortex, diencephalon including the hypothalamus and thalamus, hippocampus, autonomic ganglia, sensory fibers, pituitary gland including neurohypophysis.	
Neuroendocrine changes seem to undergo change over increased time of exposure. Neurosecretion in the hypothalamus and in the pituitary each go through a complex sequence over time, where EMF exposure initially produces increased hormone secretion but where over time, the neurosecretory cells become "exhausted", leading to lowered secretion and in some cases cell death (pp. 77–96).	Elevated [Ca2+]i stimulates hormone secretion. However when such elevated [Ca2+]i occurs over extended time periods it is highly damaging to the cell, leading in some cases to apoptosis; thus this time course of action should not be surprising.
Histological changes include boutons/argyrophilia, smaller neurons, vacuole formation in neuroendocrine cells, bead-like thickening along dendrites (pp. 66, 70, 71, 73, 97, 98, 100, 111, 115–117, 121–125). Spines near the ends of dendrites become deformed and with still more sessions of irradiation, disappeared entirely (p. 70). Sensory neurons, following exposures, developed changes characteristic of irritation, with "marked tortuosity of the nerve fibers." Many histological changes are seen in the hypothalamic cells (pp. 87–92) as their neuroendocrine function becomes impacted. Histological changes were found even with exposures that produced no apparent functional changes.	
Many histological and functional changes are reported to initially be reversible, following cessation of exposure, but progressively become irreversible with longer exposure. (pp. 64, 72, 74). Paralleling the development of irreversibility, it is found that "Repeated exposure leads to gradual increase in severity of observed changes." including "increasingly severe disturbance of conditioned reflex activity in the animals, changes in responses of animals particularly sensitive to acoustic stimulation" (p. 104).	If this is also true in humans, then claims that there cannot be non-thermal effects, claims which act to prolong exposures, may be causing irreversible damage to many humans.
EEG changes (pp. 55, 60, 102), including seizure activity following sensory provocation.	Lai (1997) has an extensive review of EEG changes in animals following non-thermal microwave EMF exposures
Synaptic connections in regions of the brain are disrupted (pp. 65–74, 97, 113, 121, 136), and at the extreme, some neurons are completely asynaptic (p. 73).	Synaptic connections are known to be disrupted in autism; could this suggest that autism may be generated by EMF exposure? No doubt, we need much more evidence on this.
"after prolonged and repeated irradiation with low-intensity centimeter waves, with no elevation of the body temperature and when the animal's condition remained satisfactory, changes were nevertheless found in the sensory fibers of the skin and viscera in the form of irritation phenomena. These findings concur with the view in the literature that the receptor system as a whole and, in particular its preterminal portions are highly sensitive." p. 76. This description is similar to what is reported to occur in electromagnetic	One wonders whether almost 60 years ago, the Soviet literature may have already described a possible animal model for EHS. None is known to exist today, and because of that, EHS studies are severely constrained. Clearly one needs to be skeptical about this interpretation, but it is of great importance that this be further studied.

neuropsychiatric effects are produced by exposure to various nonthermal microwave frequency EMFs. Perhaps the most important of these 26 is the Santini et al. (2003) study of people living near cell phone base stations.

hypersensitivity (EHS). Other such studies are described and include cumulative changes over time, that may also explain changes reported in EHS (pp. 75, 99, 100, 104).

There are three recent studies on the generation of headache during or shortly following long mobile phone calls (listed under Chu et al., 2011 in Table 3). The timing of development of these headaches and the finding that they occur on the ipsilateral side of the head, the side receiving much higher EMF exposure during the call, both argue strongly that these headaches are caused by the long mobile phone calls. Such causality was concluded earlier by Frey (1998) based on earlier studies and is now still more strongly documented.

#### 7. Criteria for assessing causality in epidemiological studies

It is important to consider the different criteria that allow one to judge whether a cause and effect relationship is justified by the studies listed in Table 3 and the individual studies cited in Raines (1981). There are five such criteria that should be considered in making that judgment (see pp. 39–43 in Hennekens and Buring, 1989):

Strength of Association: Is there a strong correlation between exposure and the neuropsychiatric symptoms? There clearly is for several studies cited in Raines (1981). One example is the Dwyer and Leeper (1978) study (see Table 3) where there is a large increase in symptoms and where that increase is greater with longer occupational exposure. Another example is the Lerner (1980) study of 1300 microwave workers, where workers with relatively low exposure levels had an approximate doubling of neurological complaints and where those with substantially higher exposure levels had an approximate tripling of neurological complaints over controls. Sadcikova (1974) found that 7 of 8 neuropsychiatric symptoms studied, showed a statistically significant rise in prevalence with longer occupational exposure (see Table 3). Sadcikova (1974), also found that microwave workers had increases of 3 to over 10-fold in: feeling of heaviness in the head; tiredness; irritability; sleepiness; partial loss of memory; and skin sensitivity. There is also a strong association where important new exposures occur - this is clearly the case with all of the studies of people living near cell/mobile phone base

#### Table 3

Neuropsychiatric symptoms apparently produced by exposure to various electromagnetic fields.

Citation	EMF exposure	Apparent neuropsychiatric symptoms
Abdel-Rassoul et al. (2007)	Living near mobile phone base station	Significant increases in neuropsychiatric complaints included: headache, memory changes, dizziness, tremors, depressive symptoms, sleep disturbance; attributed to
Al-Khlaiwi and Meo (2004)	Mobile phone use	effects of EMFs on the human nervous system. Higher prevalence of fatigue, headache, dizziness, tension and sleep disturbance: the authors conclude that mobile phone use is a risk factor for developing these symptoms.
Altpeter et al. (2000)	Short-wave broadcasting tower, ranging from 6.1 to 21.8 MHz	Sleep disruption shown to occur, correlated with exposures and apparent increase over time; short term suppression of melatonin shown, based on melatonin increases during a 3 day period when the tower was turned off
Bortkiewicz et al. (2004)	Living near cell phone base station EMFs	Sleep disturbance, irritability, depression, blurred vision, concentration difficulties, nausea, lack of appetite, headache, vertigo.
Bortkiewicz et al. (2012)	Living near mobile phone base stations	Dose response relationships for sleep disturbance, irritability, depression, blurred vision, concentration difficulties, nausea, lack of appetite.
Chu et al. (2011), also Chia et al. (2000), Oftedal	Mobile phone use	Headache during prolonged mobile phone use or within an hour following such use, with pain occurring on the ipsilateral side of the head; similar observations
et al. (2000) Conrad (2013)	Smart meter EMF exposure	obtained in each of the 3 studies in column 1; see also Frey (1998). 14 common new symptoms (both severe and moderate) among those exposed and symptomatic, 13 apparent neuropsychiatric: Insomnia, tinnitus, pressure in the head, concentration difficulty, headaches, memory problems, agitation, dizziness, fatigue, skin tingling/burning, involuntary muscle contractions, eye/vision problems, numbness; These ranged in prevalence from 63% to 19% of those experiencing symptoms, such that most symptomatic people experienced multiple symptoms.
Dasdag et al. (1992)	People working in MW broadcasting or at a television transmitter station	These groups suffered from headache, fatigue, irritability, stress, sleepiness, loss of appetite, loss of hearing.
Dwyer and Leeper (1978)	People working in radiofrequency EMFs	Headache, eyestrain, dizziness, disturbed sleep, daytime sleepiness, moodiness, mental depression, memory impairment, muscle and/or cardiac pain, breathing difficulties, increased perspiration, difficulty with sex life.
Eger and Jahn (2010)	Living near mobile phone base station	Neuropsychiatric symptoms, with most showing dose-response relationships: depression; headache; cerebral symptoms; dizziness; disorders of optical and acoustic sensory systems; sleep disturbance; skin changes; with the exception of dizziness, all of these had $p < 0.001$ .
Johnson Liakouris (1998)	Study of personnel in U.S. embassy in Moscow exposed to microwave EMFs	Statistically significant increases in neurological (peripheral nerves and ganglia), dermographism (skin responses), irritability, depression, loss of appetite, concentration difficulties, peripheral ganglia and nerve dysfunction.
Khan (2008)	Excessive mobile phone use	Complaints of headache, fatigue, impaired concentration, memory disturbance, sleeplessness, hearing problems.
Kolodynskii and Kolodinska (1996)	Children living near a Radio Location Station, Latvia	Memory dysfunction, attention dysfunction, lowered motor function, slowed reaction time, lowered neuromuscular endurance.
Lamech (2014)	Exposure to wireless smart meter radiation in Victoria, Australia	The most frequent symptoms to develop after smart meter radiation exposure were insomnia, headache, tinnitus, fatigue, cognitive disturbances, dysesthesias (abnormal sensation), dizziness.
Navarro et al. (2003)	Living near cell phone base station	Statistically significant dose response relationships for fatigue, irritability, headache, nausea, loss of appetite, sleep disorder, depressive tendency, feeling of discomfort, difficulty of concentration, loss of memory, visual disorder & dizziness.
Oberfeld et al. (2004)	Living near cell phone base station	Statistically significant dose-response relationships for headache, fatigue, irritability, loss of appetite, visual disorder, nausea, sleeping disorders, dizziness, poor concentration, memory loss.
Oto et al. (1994)	Occupational exposure of 25 workers to either UHF television broadcasting (10) or to 1062 kHz medium wave broadcasting (15)	10 neuropsychiatric changes were assessed, all showing statistically significant changes compared with controls: Somatization*, obsessive compulsivity*, interpersonal sensitivity, depression, anxiety*, hostility*, phobic anxiety*, paranoid ideation, psychoticism*, sleeping disturbance. *n < 0.001.
Sadcikova (1974)	Occupational exposure to microwave radiation, including at $< 0.7 \text{ mW/cm}^2$	Heaviness in head*, fatigue*, irritability*, sleepiness, memory loss*, cardiac pain*, dermographism (skin sensitivity)*, hyperhidrosis* * similicant increase with time of exposure
Salama and Abou El Naga (2004)	High cell (mobile) phone use	Most common effects were headache, ear ache, sense of fatigue, sleep disturbance, concentration difficulty, face burning sensation. The first three of these had very high statistical significance for correlation with extent of cell phone use.
Santini et al. (2003)	Living near cell phone base stations	Each of the following neuropsychiatric symptoms showed statistical significant dose-response relationships: nausea, loss of appetite, visual disturbance, irritability, depressive tendencies, lowered libido, headache, sleep disturbance, feeling of discomfort, fatigue.
Schüz et al. (2009)	Mobile phone use	Found a small, statistically significant increase in migraine and vertigo. Also found an apparent lowered occurrence of Alzheimer's, other dementia, Parkinson's and cpilepsy – these latter were interpreted as being due to perhaps early symptoms of the developing diseases lowering probability of acquiring a mobile phone.
Söderqvist et al. (2008)	Use of mobile phone among adolescents	Increased mobile phone use was associated with increases in tiredness, stress, headache, anxiety, concentration difficulties and sleep disturbances.
Thomée et al. (2011)	High mobile phone use	High mobile phone use was associated with statistically significant rises in stress and sleep disturbance, with somewhat weaker association with depression.
Waldmann-Selsam et al. (2009)	Digital TV signaling	Constant headaches, pressure in head, drowsiness, sleep problems, tightness in chest, shortness of breadth, depressive mood, total apathy, loss of empathy, burning skin, inner burning, leg weakness, pain in limbs, stabbing pain in various organs, weight increase.

• .

stations, listed in Table 3 and also with the two studies of people who become exposed to radiation from smart meters. The studies listed in Table 3 under Chu et al. (2011) (see also Chia et al., 2000; Oftedal et al., 2000) are of a special type. Here people making very long (over 1 h) cell/mobile phone calls develop headaches an hour or more following the initiation of the long call. So these occur within a specific time range following initiation of these long calls, such that headache would only occur very infrequently in that time frame by chance. So here again, there is a strong association. While there is no question that many of these studies show high strength of association, it is also clear that it is becoming progressively more difficult to do these studies. As exposures become almost universal in countries around the world, it is getting difficult if not impossible to find good negative controls. There may be a similar problem in doing animal studies, such that it may be necessary to raise animals in Faraday cages in order to avoid exposures that would otherwise occur as a consequence of our near ubiquitous EMFs.

*Biological credibility* is extremely strong here, with three aspects of the biology predicting that these low intensity fields cause widespread neuropsychiatric effects. This was discussed above and is reconsidered in the following section.

Consistency within the different epidemiological studies and with other types of studies. The epidemiological studies listed in Table 3 and also those showing neuropsychiatric effects that were cited in Raines (1981) have been performed in many different countries with different cultures. They have been performed in multiple countries in Western Europe, Eastern Europe, the Middle East and in East Asia, as well as in the U.S. and Australia. They are, therefore, not limited to one or two cultural contexts. This is deemed, therefore, an important indicator of causality. We also have a surprising consistency of apparent neuropsychiatric effects of different fields, including various occupational exposures and exposures to cell/mobile phone base stations, exposure to the phones themselves, exposure to smart meter pulses, and other EMFs (see Table 3). Pulsation patterns, frequencies and exact intensities may produce various biological responses (Pall, 2015; Pangopoulos et al., 2013; Belyaev, 2015) so it is a bit surprising that we have as much consistency as we do have across different types of exposures. We also have consistency with the biology discussed in the previous section. Because elevated VGCC activity produced by genetic polymorphism (Table 1) produces diverse neuropsychiatric effects, it is not surprising that elevation of VGCC activity produced by microwave EMF exposure apparently also produces diverse neuropsychiatric effects. Similarly because non-thermal EMF exposures produce widespread changes in brain structure and function in animals (Tolgskaya and Gordon, 1973), it is not surprising that the neuropsychiatric symptoms, which are produced as a consequence of brain dysfunction are produced by such EMFs.

Time sequence: It is clear that the all of these effects follow exposure in the various studies that have been published. In some studies, it is also clear that longer occupational exposure times produce increased symptom prevalence. These include Dwyer and Leeper (1978) and Baranski and Edelwejn (1975). These observations all support a causal relationship between exposure to EMF and the development of neuropsychiatric symptoms.

Dose-response relationship: It is assumed, here, that biological effects have a positive correlation with the intensity of the apparent causal stressor. This is not necessarily true of EMF effects, because it has been shown that there are "window effects" where specific intensities have larger biological effects, than do either lower or higher intensities (Pall, 2015; Pangopoulos et al., 2013; Belyaev, 2015). Nevertheless, where different intensities were studied in these epidemiological studies, they do show the dose-response relationship assumed here including Altpeter et al.

(2000), Dwyer and Leeper (1978), Eger and Jahn (2010), Lerner (1980), Navarro et al. (2003), Oberfeld et al. (2004), Salama and Abou El Naga (2004), Santini et al. (2003) and Thomée et al. (2011). Thus these data do fit well to the assumed dose-response relationship, found in most causal roles. The Altpeter et al. (2000) study showed a special type of evidence for causality: during a 3day period when the broadcasting tower was turned off, the melatonin levels recovered to near-normal levels. The studies of headache occurrence on prolonged cell/mobile phone calls (typically well over one hour) listed under Chu et al. (2011) in Table 3 also suggest the assumed dose-response relationship (see also Chia et al., 2000; Oftedal et al., 2000 and earlier citations listed in Frey, 1998). Because such headaches only occur with prolonged cell/mobile phone calls, these studies also provide evidence for a dose-response relationship because low doses are ineffective. Furthermore these same studies provide evidence for such a doseresponse relationship from another type of observation. Because the headaches occur predominantly on the ipsilateral side of the head which receives much higher EMF exposure intensity, rather than on the contralateral side of the head, which receives much lower intensities, this provides an additional type of evidence for the predicted dose-response relationship.

While the evidence is convincing that the various neuropsychiatric apparent consequences of microwave EMF exposure are in fact caused by such exposures, there may be somewhat more controversy about another EMF-neuropsychiatric linkage. Havas et al. (2010) have reported a similar list of neuropsychiatric symptoms in electromagnetic hypersensitivity (EHS) patients. They found that each of the following symptoms were common in EHS: poor short term memory; difficulty of concentration; eye problems; sleep disorder; feeling unwell; headache; dizziness; tinnitus; chronic fatigue: tremors: body pain: difficulty speaking: tingling sensation in feet or hands; difficulty writing; difficulty walking; migraine. The similarity of these symptoms to the most commonly found symptoms following non-thermal microwave EMF exposures (Table 3), suggests that EHS is a genuine sensitivity to EMFs. In the bottom row in Table 2, sensitivities were found in rodent studies following non-thermal exposure that suggest a possible animal model for the study of EHS. Each of these EHS-related issues needs to be followed up experimentally.

#### 8. Discussion and conclusions

In the previous section, each of the five criteria for assessing whether an epidemiological association is causal, were considered. Those five are (Hennekens and Buring, 1989): (1) strength of association; (2) biological credibility; (3) consistency; (4) time sequence; (5) dose-response relationship. Each of these five provide strong support for causality such that the combination of all five provides compelling evidence for causality. Low-intensity microwave frequency EMFs do cause diverse neuropsychiatric symptoms. While each of these five is important here, the one that is most important is the criterion of biological credibility.

Three related sets of biological observations each predict that low-intensity microwave EMFs produce widespread neuropsychiatric effects:

 Such EMFs act via activation of VGCCs, acting through the VGCC voltage sensor which is predicted to be exquisitely sensitive to these EMFs (Pall, 2015). VGCCs occur in high densities throughout the nervous system and have essential roles throughout the nervous system in releasing neurotransmitters and neuroendocrine hormones. These properties predict, therefore, that these low intensity non-thermal microwave EMFs cause widespread changes in the nervous system, causing, in turn, diverse neuropsychiatric effects.

- Elevated VGCC activity, produced by an allele of the CACNA1C gene which encodes the channel of the main L-type VGCC in the brain, produces various neuropsychiatric effects (Table 1). This predicts, that low intensity non-thermal microwave frequency EMFs which also produce elevated L-type and other VGCC activity, therefore produce widespread neuropsychiatric effects.
- 3. Studies reviewed in the Tolgskaya and Gordon, 1973 publication (Table 2) have shown that the cells of the mammalian nervous system show high sensitivity to various non-thermal microwave and lower frequency EMFs, being apparently more sensitive than any other organ in the body of rodents. These studies predict that the human nervous system is likely to be similarly sensitive to these EMFs, predicting, therefore, widespread neuropsychiatric effects in humans.

We not only have biological credibility but also more importantly, each of these distinct but interrelated biological considerations predicts that low-intensity, non-thermal microwave EMFs produce widespread neuropsychiatric effects. That common prediction is verified by extensive data summarized in citations provided by the Naval Medical Research Institute Research Report (June 1971), data provided by The Raines (1981) NASA report, and by 26 epidemiological studies summarized in Table 3.

The most commonly reported neuropsychiatric symptoms from these studies are summarized in Table 4.

A total of 22 different studies described in Table 3 were used for data for this table, but not 4 others that only assessed a single neuropsychiatric end point. The Altpeter study which only assessed sleep disturbance/melatonin depletion and the three studies listed under Chu et al. which only assessed headache occurrence following long cell phone calls, listed in Table 3 were not included. Because many of the studies only assessed from 3 to 7 specific symptoms, it is not surprising that the numbers of studies reporting a specific symptom fall far below 22. Where several symptom descriptions were included under one heading, such as dysesthesia, if a study had more than one of these symptom descriptions, it was only counted once.

All the symptoms listed in Table 4 should be considered established parts of microwave syndrome (Hocking, 2001; Johnson Liakouris, 1998). Even if the statistical significance in each study was of the lowest statistical significance (p < .05) one would expect only 1 positive study to occur at random out of the 22 studies included here. Because many individual symptoms were not surveyed in many individual studies, the expectation is

#### Table 4

Commonly reported neuropsychiatric symptoms following microwave EMF exposure.

Symptom(s)	Numbers of studies reporting
Sleep disturbance/insomnia	17
Headache	14
Fatigue/tiredness	11
Depression/depressive symptoms	10
Dysesthesia (vision/hearing/olfactory dysfunction)	10 .
Concentration/attention/cognitive dysfunction	10
Dizziness/vertigo	9
Memory changes	8
Restlessness/tension/anxiety/stress/ agitation/feeling of discomfort	8
Irritability	7
Loss of appetite/body weight	6
Skin tingling/burning/inflammation/ dermographism	6
Nausea	5

substantially lower than that. Each of these, having shown positive results in 5 or more studies are highly unlikely, therefore, to have occurred by chance. Stong statistical significance is also seen for individual neuropsychiatric effects reported to have p < 0.001 in the Eger and Jahn (2010) and Oto et al. (1994) studies (see Table 3).

EEG changes may well be part of microwave syndrome, as well. While none of the studies described in Table 3 measured EEGs, six studies of human occupational exposure cited in the Raines (1981) showed EEG changes (Baranski and Edelwein, 1975; Bise, 1978; Dumanskij and Shandala, 1974; Lerner, 1980; Sheppard and Eisenbud, 1977). Murbach et al. (2014) cited 10 human studies in support of their statement that "the most consistently reported effects (of mobile phone use) in various studies conducted by different laboratories are changes in the electroencephalogram (EEG) power spectrum." Three recent studies (Lustenberger et al., 2013; Schmid et al., 2012a,b) and several earlier studies cited in Wagner et al. (1998) have each shown EEG changes in sleeping humans exposed to non-thermal pulsed microwave fields. Two recent studies showed EEG changes in persons exposed to Wi-Fi fields (Maganioti et al., 2010; Papageorgiou et al., 2011). Lai (1997) described 8 animal studies showing changes in EEG patterns in animals exposed to non-thermal EMFs and three additional animal studies were described in Tolgskaya and Gordon (1973). With the exception of the 6 studies cited in the second sentence in this paragraph, all of these are direct experimental studies which are not, therefore, susceptible to the questions of causality that can be raised about epidemiological studies. It is the author's view that future studies should consider studying EEG changes as an objectively measurable assessment of brain physiology and that before and after increased exposure studies should be considered when a new EMF source is to be introduced into human populations. While such studies must be done carefully, given the complexity of EEGs, even very small numbers of individuals may produce highly statistically significant results in well designed studies analyzed with paired t-tests.

One of the citations from the previous paragraph, Bise (1978) reviewed earlier studies of low level microwave frequency exposures in humans and concluded that such EMFs produced the following neuropsychiatric effects: headache, fatigue, irritability, dizziness, loss of appetite, sleepiness, sweating, difficulty of concentration, memory loss, depression, emotional instability, dermographism, tremor, hallucinations and insomnia. The strong similarity of this list from 37 years ago and the list in Table 4 should be noted. The Bise (1978) list is based on occupational exposure studies whereas the current list in Table 4 is based primarily on EMF exposures from cell/mobile phone base stations, from heavy cell phone usage and from smart meters, three types of exposures that did not exist in 1978. The strong similarity between the Bise (1978) list and the current one 37 years later alone produces a compelling argument that the 11 neuropsychiatric effects found on both lists are caused by exposure to multiple types of low-intensity microwave EMFs.

The pattern of evidence is compelling in support of the earlier statement of Levitt and Lai (2010) that "the primary questions now involve specific exposure parameters, not the reality of complaints or attempts to attribute such complaints to psychosomatic causes, malingering or beliefs in paranormal phenomena."

We can barely imagine how the combinations of neuropsychiatric effects, including those in Table 4, will influence human behavior and social interactions, now that the majority of the human populations on earth are exposed to ever increasing intensities and diversity of microwave frequency EMFs. You may recall that three of the occupational exposure studies cited in (Raines, 1981 showed increasing prevalence of neuropsychiatric symptoms with years of exposure to consistent patterns of EMF exposure intensities (Dwyer and Leeper, 1978; Sadcikova, 1974; Baranski and Edelwein, 1975). With ever increasing exposures in human populations, we have no idea what the consequences of these ever increasing exposures will be.

#### **Conflict of interest**

The author declares no conflict of interest.

#### References

- Abdel-Rassoul, G., El-Fateh, O.A., Salem, M.A., Michael, A., Farahat, F., El-Batanouny, M.A., Salem, F., 2007, Neurobehavioral effects among inhabitants around mobile phone stations. Neurotoxicology 28, 434-440.
- Adey, W.R., 1993. Biological effects of electromagnetic fields. J. Cell. Biochem. 51, 410-416.
- Al-Khlaiwi, T., Meo, S.A., 2004. Association of mobile phone radiation with fatigue, headache, dizziness, tension and sleep disturbance in Saudi population. Saudi Med. J. 25, 732-736.
- Altpeter, E., Battaglia, M., Bader, A., Pluger, D., Minder, C.E., Abelin, T., 2000. Ten Years Experience with Epidemiological Research in the Vicinity of the Short-Wave Broadcasting Area Schwarzenburg: What does the Story Tell Us? , http://www.salzburg.gv.at/Proceedings\_%2819%29\_Altpeter.pdf.
- Ammari, M., Brillaud, E., Gamez, C., Lecomte, A., Sakly, M., Abdelmelek, H., de Seze, R., 2008a. Effect of a chronic GSM 900 MHz exposure on glia in the
- rat brain. Biomed. Pharmacother. 62, 273–281. Ammari, M., Lecomte, A., Sakly, M., Abdelmelek, H., de-Seze, R., 2008b. Exposure to GSM 900 MHz electromagnetic fields affects cerebral cytochrome c oxidase activity. Toxicology 250, 70–74. Baranski, S., Edelwejn, Z., 1975. Experimental morphologic and
- electroencephalographic studies of microwave effects on the nervous system. Ann. N. Y. Acad. Sci. 47, 109-116.
- Bas, O., Odaci, E., Kaplan, S., Acer, N., Ucok, K., Colakoglu, S., 2009. 900 MHz electromagnetic field exposure affects qualitative and quantitative features of hippocampal pyramidal cells in the adult female rat. Brain Res. 1265, 178-185
- Belyaev, 1, 2015. Biophysical mechanisms for nonthermal microwave effects. In: Markov, Marko S. (Ed.), Electromagnetic Fields in Biology and Medicine. CRC Press, New York, pp. 49-67.
- Berridge, M.J., 1998. Neuronal calcium signaling. Neuron 21, 13-26.
- Bhat, S., Dao, D.T., Terrillion, C.E., Arad, M., Smith, R.J., Soldatov, N.M., Gould, T.D., 2012. CACNA1C (Cav1.2) in the pathophysiology of psychiatric disease. Prog. Neurobiol. 99, 1-14.
- Bigos, K.L., Mattay, V.S., Callicott, J.H., Straub, R.E., Vakkalanka, R., Kolachana, B., Hyde, T.M., Lipska, B.K., Kleinman, J.E., Weinberger, D.R., 2010. Genetic variation in CACNATC affects brain circuitries related to mental illness. Arch. Gen. Psychiatry 67, 939-945.
- Bise, W., 1978. Low power radio-frequency and microwave effects on human electroencephalogram and behavior. Physiol. Chem. Phys. 10, 387-398.
- Bolen, S.M., 1994. Radiofrequency/Microwave Radiation Biological Effects and Safety Standards: A Review, AD-A282 886, Rome Laboratory, U.S. Air Force Material Command, Griffiss Air Force Base, New York.
- Bortkiewicz, A., Zmyslony, M., Szyjkowska, A., Gadzicka, E., 2004. Subjective symptoms reported by people living in the vicinity of cellular phone base stations: review. Med. Pr. 55, 345-351. Bortkiewicz, A., Gadzicka, E., Szyjkowska, A., Politański, P., Mamrot, P., Szymczak,
- W., Zmyślony, M., 2012. Subjective complaints of people living near mobile phone base stations in Poland. Int. J. Occup. Med. Environ. Health 25, 31-40.
- Brillaud, E., Piotrowski, A., de Seze, R., 2007. Effect of an acute 900 MHz GSM exposure on glia in the rat brain: a time-dependent study. Toxicology 238, 23-33.
- Carballo-Quintás, M1, Martínez-Silva, I., Cadarso-Suárez, C., Alvarez-Figueiras, M., Ares-Pena, F.J., López-Martín, E., 2011. A study of neurotoxic biomarkers, c-fos and GFAP after acute exposure to GSM radiation at 900 MHz in the picrotoxin model of rat brains. Neurotoxicology 32, 478-494.
- Chia, S.E., Chia, H.P., Tan, J.S., 2000. Prevalence of headache among handheld cellular telephone users in Singapore: a community study. Environ. Health Perspect, 108, 1059-1062.
- Chu, M.K., Song, H.G., Kim, C., Lee, B.C., 2011, September, Clinical features of headache associated with mobile phone use: a cross-sectional study in university students. BMC Neurol. 11, 115, http://dx.doi.org/10.1186/1471-2377-11-115,
- Conrad, R.H., 2013. Smart Meter Health Effects Survey and Report, http://www. mainecoalitiontostopsmartmeters.org/wp-content/uploads/2013/02/Exhibit-D-Smart-Meter-Health-Effects-Report-w-AppendicesV3-1-9Reduced-Appendices.pdf.
- Dasdag, S., Akdag, M.Z., Ulukaya, E., Uzunlar, A.K., Ocak, A.R., 2009. Effect of mobile phone exposure on apoptotic glial cells and status of oxidative stress in rat brain. Electromagn. Biol. Med. 28, 342-354.
- Dasdag, S., Akdag, M.Z., Kizil, G., Kizil, M., Cakir, D.U., Yokus, B., 2012. Effect of 900 MHz radio frequency radiation on beta amyloid protein, protein carbonyl, and malondialdehyde in the brain. Electromagn. Biol. Med. 31, 67-74.
- Dasdag, S., Balci, K., Celik, M.S., Batun, S., Kaplan, A., Bolaman, Z., Tekes, S., Akdag, Z., 1992. Neurologic and biochemical lindings and CD4/CD8 ratio in

people occupationally exposed to RF and microwave. Biotechnol, Biotechnol. Fmin. 6 (4), 37-39

- Dumanskij, J.D., Shandala, M.G., 1974. The biologic action and hygienic significance of electromagnetic fields of super-high and ultrahigh frequencies in densely populated areas. In: Czerski, P., et al. (Eds.), Effects and Health Hazards of Microwave Radiation, Proceedings of an International Symposium, Warsaw15-18 October 1973.
- Dunlap, K., Luebke, J.L., Turner, T.J., 1995. Exocytic Ca\*\* channels in the mammalian central nervous system. Neuroscience 18, 89-98.
- Dwyer, M.J., Leeper, D.B., 1978. A Current Literature Report on the Carcinogenic Properties of Ionizing and Nonionizing Radiation. DHEW Publication (NIOSH), , pp. 78+134.
- Eberhardt, J.L., Persson, B.R., Brun, A.E., Salford, L.G., Malmgren, L.O., 2008. Blood-brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones. Electromagn. Biol. Med. 27, 215-229.
- Eger, H., Jahn, M., 2010. Specific symptoms and radiation from mobile basis stations in Selbitz, Bavaria, Germany: evidence for dose-effect relationship. Umw. - Med. Ges. 23, 130-139.
- Frey, A.H., 1998. Headaches from cellular telephones: are they real and what are the implications? Environ. Health Perspect. 106, 101-103.
- Grafström, G., Nittby, H., Brun, A., Malmgren, L., Persson, B.R., Salford, L.G., Eberhardt, J., 2008. Histopathological examinations of rat brains after longterm exposure to GSM-900 mobile phone radiation. Brain Res. Bull. 77, 257-263.
- Havas, M., Marrongelle, J., Poliner, B., Kelley, F., Rees, C.R.G., Tully, L., 2010. Provocation study using heart rate variability shows microwave radiation from 2.4 GHz phone affects autonomic nervous system. Eur. J. Oncol. Libr. 5, 273-300
- Hennekens, C.H., Buring, J.E., 1989. Epidemiology in Medicine. Little Brown and Co., Boston.
- Hocking, B., 2001. Microwave sickness: a reappraisal. Occup. Med. 51, 66-69.
- Johnson Liakouris, A.G., 1998, Radiofrequency (RF) sickness in the Lilienfeld study: an effect of modulated microwaves? Arch. Environ. Health 53, 226-228.
- Khan, M.M., 2008. Adverse effects of excessive mobile phone use. Int. J. Occup. Med. Environ, Health 21, 289-293.
- Khurana, V.G., Hardell, L., Everaert, J., Bortkiewicz, A., Carlberg, M., Ahonen, M., 2010. Epidemiological evidence for a health risk from mobile phone base stations. Int. J. Occup. Environ. Health 16, 263-267.
- Kolodynskii, A.A., Kolodinska, V.V., 1996. Motor and psychological functions of school children living in the area of the Skrunda Radio Location Station in Latvia. Sci. Total Environ, 180, 87-93.
- Krug, A., Nieratschker, V., Markov, V., Krach, S., Jansen, A., Zerres, K., Eggermann, T., Stöcker, T., Shah, N.J., Treutlein, J., Mühleisen, T.W., Kircher, T., 2010. Neuroimage 49, 1831-1836.
- Krug, A., Witt, S.H., Backes, H., Dietsche, B., Nieratschker, V., Shah, N.J., Nöthen, M.M., Rietschel, M., Kircher, T., 2014. A genome-wide supported variant in CACNATC influences hippocampal activation during episodic memory encoding and retrieval. Eur. Arch. Psychiatry Clin. Neurosci. 264, 103-110.
- Kumlin, T., fivonen, H., Micttinen, P., Juvonen, A., van Groen, T., Puranen, L., Pitkäaho, R. Juutilainen, J., Tanila, H., 2007. Mobile phone radiation and the developing brain: behavioral and morphological effects in juvenile rats. Radiat. Res. 168, 471-479.
- Lai, H., 1997. Neurological effects of radiofrequency electromagnetic radiation relating to wireless communication technology. In: Paper presented at the IBC-UK Conference: "Mobile Phones – Is There a Health Risk?". , http:// www.papcruzin.com/radiofrequency/henry\_lai1.htm.
- Lamech, F., 2014. Self-reporting of symptom development from exposure to radiofrequency fields of wireless smart meters in Victoria, Australia: a case series. Altern. Ther. Health Med. 20, 28-39.
- Lerner, E.J., 1980, RF radiation: biological effects. IEEE Spectr. 17 (December), 51-59
- Levitt, B.B., Lai, H., 2010. Biological effects from exposure to electromagnetic radiation emitted by cell towers base stations and other antenna arrays. Environ. Rev. 18, 369-395.
- Li, Y., Yan, X. Liu, J., Li, L., Hu, X., Sun, H., Tian, J., 2014. Pulsed electromagnetic field enhances brain-derived neurotrophic factor expression through L-type voltage-gated calcium channel- and Erk-dependent signaling pathways in neonatal rat dorsal root ganglion neurons. Neurochem. Int. 75, 96-104.
- Lisi, A., Ledda, M., Rosola, E., Pozzi, D., D'Emilia, E., Giuliani, L., Foletti, A., Modesti, A., Morris, S.J., Grimaldi, S., 2006. Extremely low frequency electromagnetic field exposure promotes differentiation of pituitary corticotrope-derived AtT20 D16V cells. Bioelectromagnetics 27, 641-651.
- López-Martín, E., Relova-Quinteiro, J.L., Gallego-Gómez, R., Peleteiro-Fernández, M., Jorge-Barreiro, F.J., Ares-Pena, F.J., 2006. GSM radiation triggers seizures and increases cerebral c-Fos positivity in rats pretreated with subconvulsive doses of picrotoxin. Neurosci. Lett. 398, 139-144.
- Lustenberger, C., Murbach, M., Dürr, R., Schmid, M.R., Kuster, N., Achermann, P., Huber, R., 2013. Stimulation of the brain with radiofrequency electromagnetic field pulses affects sleep-dependent performance improvement. Brain Stimul. 6, 805-811.
- Maganioti, A.E., Papageorgiou, C.C., Hountala, C.D., Kyprianou, M.A., Rabavilas, A.D., Papadimitriou, G.N., Capsalis, C.N., 2010. Wi-Fi electromagnetic fields exert gender related alterations of EEG. In: 6th International Workshop on

2

 $Biological \ Effects \ of \ Electromagnetic \ Fields, \ Bodrun, \ Turkey, \ October, \ http://www.istanbul.edu.tr/6internatwshopbioeffemf/.$ 

- Mausset-Bonnefont, A.L., Hirbec, H., Bonnefont, X., Privat, A., Vignon, J., de Seze, R., 2004. Acute exposure to GSM 900 MHz electromagnetic fields induces glial reactivity and biochemical modifications in the rat brain. Neurobiol. Dis. 17, 445-454.
- Murbach, M., Neufeld, E., Christopoulou, M., Achermann, P., Kuster, N., 2014. Modeling of EEG electrode artifacts and thermal ripples in human radiofrequency exposure studies. Bioelectromagnetics 35, 273–283.
- Naval Medical Research Institute Research Report, 1971, June. Biobliography of Reported Biological Phenomena ("Effects") and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised.
- Navarro, G., Segure, J., Porteles, M., Perretta, Gomez, 2003. The microwave syndrome: study in Spain. Electromag. Biol. Med. 22, 161–169.
- Oberfeld, G., Navarro, A.E., Portoles, M., Maestu, C., Gomez-Perretta, C., 2004. The microwave syndrome: further aspects of a Spanish study, http://www.apdr. info/electrocontaminacion/Documentos/Investigacion/ESTUDOS%20 EPIDEMIOLOXIDOS%20E%20ANTENAS/The%20Microwave%20Syndrome%20-% 20Further%20Aspects%20of%20a%20Spanish%20Study.pdf.
- Odaci, E., Bas, O., Kaplan, S., 2008. Effects of prenatal exposure to a 900 MHz electromagnetic field on the dentate gyrus of rats: a stereological and histopathological study. Brain Res. 1238, 224–249.
- Oftedal, G., Wilén, J., Sandström, M., Mild, K.H., 2000. Symptoms experienced in connection with mobile phone use. Occup. Med. (Lond.) 50, 237–245.
- Oto, R., Akdag, Z., Dasdag, S., Celik, Y., 1994. Evaluation of psychologic parameters in people occupationally exposed to radiofrequencies and microwave. Biotechnol. Biotechnol. Equip. 8 (4), 71–74.
- Pall, M.L., 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. J. Cell. Mol. Med. 17, 958–965.
- Pall, M.L., 2014. Electromagnetic field activation of voltage-gated calcium channels: role in therapeutic effects. Electromagn. Biol. Med. 33, 251.
- Pall, M.L., 2015. Review: scientific evidence contradicts findings and assumptions of Canadian safety panel 6: microwaves act through voltagegated calcium channel activation to induce biological impacts a non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. Rev. Environ. Health 30, 99–116.
- Pangopoulos, D.J., Johansson, O., Carlo, G.L., 2013. Evaluation of specific absorption rate as a dosimetric quantity for electromagnetic fields bioeffects. PLOS ONE 8 (6), e62663.
- Papageorgiou, C.C., Hountala, C.D., Maganioti, A.E., Kyprianou, M.A., Rabavilas, A.D., Papadimitriou, G.N., Capsalis, C.N., 2011. Effects of wi-fi signals on the p300 component of event-related potentials during an auditory hayling task. J. Integr. Neurosci. 10, 189–202.
- Pilla, A.A., 2012. Electromagnetic fields instantaneously modulate nitric oxide signaling in challenged biological systems. Biochem. Biophys. Res. Commun. 426, 330–333.
- Rağbetli, MC1, Aydinlioğlu, A., Koyun, N., Rağbetli, C., Bektas, S., Ozdemir, S., 2010. The effect of mobile phone on the number of Purkinje cells: a stereological study. Int. J. Radiat. Biol. 86, 548–554.
- stereological study. Int. J. Radial. Biol. 86, 548-554.
  Raines, J.K., 1981, April. Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories. National Aeronautics and Space Administration, NASA CR 166661, Greenbelt, Maryland.
- Sadcikova, M.N., 1974. Clinical manifestations of reactions to microwave radiation in various occupational groups. In: Czerski, P. (Ed.), Biological Effects and Health Hazards of Microwave Radiation. Proceedings of the International Symposium, Warsaw 13–18 October 1973. Polish Med Publishers, Warsaw, pp. 261–267.
- Salama, O.E., Abou El Naga, R.M., 2004. Cellular phones: are they detrimental? J. Egypt. Public Health Assoc. 79, 197–223.

- Salford, L.G., Brun, A.E., Eberhardt, J.L., Malmgren, L., Persson, B.R., 2003. Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. Environ. Health Perspect. 111, 881–883.
- Santini, R., Santini, P., Le Ruz, P., Danze, J.M., Seigne, M., 2003. Survey of people living in the vicinity of cellular phone base stations. Electromagn. Biol. Med. 22, 41-49.
- Schmid, M.R., Loughran, S.P., Regel, S.J., Murbach, M., Bratic Grunauer, A., Rusterholz, T., Bersagliere, A., Kuster, N., Achermann, P., 2012a. Sleep EEG alterations: effects of different pulse-modulated radio frequency electromagnetic fields. J. Sleep Res. 21, 50–58.
- Schmid, M.R., Murbach, M., Lustenberger, C., Maire, M., Kuster, N., Achermann, P., Loughran, S.P., 2012b. Sleep EEG alterations: effects of pulsed magnetic fields versus pulse-modulated radio frequency electromagnetic fields. J. Sleep Res. 21, 620–629.
- Schüz, J., Waldemar, G., Olsen, J.H., Johansen, C., 2009. Risks for central nervous system diseases among mobile phone subscribers: a Danish retrospective cohort study. PLoS ONE 4 (2), e4389, http://dx.doi.org/10.1371/ journal.pone.0004389.
- Sheppard, A.R., Eisenbud, M., 1977. Biological Effects of Electric and Magnetic Fields of Extremely Low Frequency. New York University Press, New York
- Fields of Extremely Low Frequency. New York University Press, New York. Söderqvist, F., Carlberg, M., Hardell, L., 2008. Use of wireless telephones and self-reported health symptoms: a population-based study among Swedish adolescents aged 15–19 years. Environ. Health 7 (May), 18, http://dx.doi.org/ 10.1186/1476-069X-7-18.
- Soeiro-de-Souza, M.G., Otaduy, M.C., Dias, C.Z., Bio, D.S., Machado-Vieira, R., Moreno, R.A., 2012. The impact of the CACNA1C risk allele on limbic structures and facial emotions recognition in bipolar disorder subjects and healthy controls. J. Affect, Disord. 141, 94–101.
- Sonmez, O.F., Odaci, E., Bas, O., Kaplan, S., 2010. Purkinje cell number decreases in the adult female rat cerebellum following exposure to 900 MHz electromagnetic field. Brain Res. 1356, 95–101.
- Tesli, M., Skatun, K.C., Ousdal, O.T., Brown, A.A., Thoresen, C., Agartz, I., Melle, I., Djutovic, S., Jensen, J., Andreassen, O.A., 2013. CACNA1C risk variant and amygdala activity in bipolar disorder. schizophrenia and healthy controls. PLOS ONE 8 (2), e56970, http://dx.doi.org/10.1371/journal.pone.0055970.
- Thimm, M., Kircher, T., Kellermann, T., Markov, V., Krach, S., Jansen, A., Zerres, K., Eggermann, T., Stöcker, T., Shah, N.J., Nöthen, M.M., Rietschel, M., Witt, S.H., Mathiak, K., Krug, A., 2011. Effects of a CACNA1C genotype on attention networks in healthy individuals. Psychol. Med, 41, 1551–1561.
  Thomée, S., Härenstam, A., Hagberg, M., 2011. Mobile phone use and stress,
- Thomée, S., Härenstam, A., Hagberg, M., 2011. Mobile phone use and stress, sleep disturbances, and symptoms of depression among young adults – a prospective cohort study. BMC Public Health 11 (January), 66, http:// dx.doi.org/10.1186/1471-2458-11-66.
- Tolgskaya, M.S., Gordon, Z.V., 1973. Pathological Effects of Radio Waves. (B. Haigh, Trans.)Consultants Bureau, New York/London, pp. 146.
- Wagner, P., Röschke, J., Mann, K., Hiller, W., Frank, C., 1998. Human sleep under the influence of pulsed radiofrequency electromagnetic fields: a polysomnographic study using standardized conditions. Bioelectromagnetics 19, 199-202.
- Waldmann-Selsam, C., Aschermann, C., Kern, M., 2009. Warning against adverse health effects from the operation of digital broadcast television stations (DVB-T): letter from 3 German physicians to the U.S. President and Congress, http://www.stayontruth.com/ internationality.com/
- warning-against-adverse-health-effects-digital/IV.php.
- Walleczek, J., 1992. Electromagnetic field effects on cells of the immune system: the role of calcium signaling, FASEB J. 6, 3177–3185.
- Wheeler, D.B., Randall, A., Tsien, R.W., 1994. Roles of N-type and Q-type channels in supporting hippocampal synaptic transmission. Science 264, 107-111.

## BEFORE THE PENNSYLVANIA PUBLIC UTILITY COMMISSION

Catherine J. Frompovich	•	
-	•	Docket No. C-2015-2474602
	:	
v.	:	
	:	
PECO Energy Company	:	

## CERTIFICATE OF SERVICE

I, Catherine J. Frompovich, hereby certify that I have this day served a copy of my Exceptions Protest due June 13, 2017 to the following via

Certified Mail \* Return Receipt Requested to: Rosemarie Chiavetta, Secretary Pennsylvania Public Utility Company Commonwealth of Pennsylvania P.O. Box 3265 Harrisburg, PA 17105-3265 Courtesy copy emailed to OSA at <u>ra-OSA@pa.gov</u>

And by First Class U.S. Postal Service with Tracking mail to:

Administrative Law Judge Darlene D Heep Pennsylvania Public Utility Commission 801 Market Street, Ste. 4063 Philadelphia, PA 19107

Ward Smith, Esq. Shawane Lee, Esq. Thomas Carl Watson, Esq. Exelon Business Services Company LLC Legal Department 2301 Market Street, S23-1 Philadelphia, PA 19103

Dated at Ambler, Pennsylvania, June 7, 2017

arturia proch

Catherine J. Frompovich, Pro Se 23 Cavendish Drive, Ambler, PA 19002 215-653-7575



JUN 7 2017

PA PUBLIC UTILITY COMMISSION SECRETARY'S BUREAU





OUSE, PA

From CJ Frompovich 23 Cavendish Drive Ambler, PA 19002

TO

Certified Mail \* Return Receipt Requested Rosemarie Chiavetta, Secretary Pennsylvania Public Utility Commission Commonwealth of Pennsylvania P.O. Box 3265 Harrisburg, PA 17105-3265